



## **Education**

### **Herniated Disk**

#### **What is a herniated disk?**

A herniated disk is a disk that has bulged out from its proper place in your back. Disks are small, circular cushions between the bones of the spine (vertebrae). Normally, disks act as shock absorbers to cushion your vertebrae from each other as you move. A herniated disk may press on nearby nerves and cause severe pain.

#### **How does it occur?**

When a disk is damaged, the soft rubbery center of the disk squeezes out through a weak point in the hard outer layer. A disk may be damaged by:

- a fall or accident
- repeated straining of your back
- a sudden strenuous action such as lifting a heavy weight or twisting violently.

A herniated disk may also happen spontaneously without any specific injury.

#### **What are the symptoms?**

If your herniated disk is in your back, your symptoms may develop gradually or begin suddenly. Symptoms include:

- back pain
- numbness, tingling, pain, or weakness in one or both legs (this is called sciatica)
- changes in bladder and bowel habits.

Symptoms of a herniated disk in your neck may also develop gradually or suddenly. You may wake up and feel a sudden aching. Or you may have a twisted neck that you cannot straighten without extreme pain. You may also have numbness, tingling, or weakness in one or both arms.

#### **How is it diagnosed?**

Your health care provider will review your symptoms and ask about the history of your pain. Then he or she will examine your spine and test the movement and reflexes in your arms and legs. Your provider may want you to have one or more of the following tests:

- x-rays of your spine
  - magnetic resonance imaging, also called MRI (an image of your spine and herniated disk generated by sound waves)
  - CT scan (computerized x-ray images of your spine)
  - electromyography (tests of electrical activity in your muscles)
  - myelography (injection of dye into the fluid around the spinal cord that can be seen on x-rays)
  - diskography (injection of dye into a disk and x-rays taken).
-

## How is it treated?

In most cases, treatment without surgery will relieve your pain.

For a herniated disk in your back, your health care provider may recommend bed rest for 1 to 2 days. You may lie flat on your back on a firm mattress or on an ordinary bed with a stiff board under the mattress. Your provider may suggest putting a pillow under your knees when you lie on your back. You may also lie on your belly with a pillow under your chest or on your side with a pillow between your legs. Use the position that is most comfortable for you.

Other treatments your provider may recommend for your back are:

- anti-inflammatory drugs
- prescription pain relievers
- muscle relaxants
- hot or cold packs
- traction
- back massage
- physical therapy
- steroid injections into the space near the herniated disk to control pain and inflammation.

Treatment for a herniated disk in your neck may include:

- hot or cold packs
- anti-inflammatory drugs
- muscle relaxants
- prescription pain relievers
- a neck collar or neck brace to relieve muscle spasms
- neck and shoulder massage
- traction, which is the process of putting bones or muscles under tension with a system of weights and pulleys to keep them from moving or to relieve pressure on them.

As your pain lessens, your health care provider will want you to begin a physical therapy program in which you will do exercises to strengthen your back muscles and joints. Stabilization exercises are also used to treat herniated disks. This therapy involves learning how to control the movement of your spine in all recreation and work activities.

If you continue to have symptoms, you may need to have surgery. However, most people who have herniated disks do not need surgery.

## How long will the effects of a herniated disk last?

The initial intense pain should go away within a few weeks, but some pain may remain for a few months. You may be prone to backaches throughout your life and therefore must remember to protect your spine when lifting or being physically active.

If the weakness and numbness in your legs continue or if you lose control of your bowel or bladder function, contact your health care provider immediately.

## How can I take care of myself?

Practice correct posture when you are walking, sitting, standing, lying down, or working.

- When lifting heavy objects, don't bend over from your waist. Kneel or squat down by the object, while keeping your back as straight as possible. Use your thigh muscles to do the lifting. Avoid twisting.
  - When you stand, always stand up straight with your shoulders back, abdomen in, and the small of the back flat. When standing for long periods, move around frequently and shift your weight from one foot to another while standing as straight as possible.
  - When you sit, have your feet flat on the floor or elevated. Get up every 20 minutes or so and stretch. Sit in a chair that has good back support.
  - Sleep on a firm mattress or one with a bed board under it. Lie on your side with your knees bent or on your back with a small pillow under your head and another pillow under your knees.
-

## What can be done to help prevent a herniated disk?

Herniated disks can often be prevented by keeping your weight down, eating a proper diet, and exercising to keep your muscles firm. Strong, flexible muscles can stabilize your spine and protect it from injury. This includes keeping your stomach muscles strong. Walking and swimming are two good exercises for strengthening and protecting your spine.

Adult Health Advisor 2006.4; Copyright © 2006 McKesson Corporation and/or one of its subsidiaries. All Rights Reserved. Developed by McKesson Provider Technologies. This content is reviewed periodically and is subject to change as new health information becomes available. The information is intended to inform and educate and is not a replacement for medical evaluation, advice, diagnosis or treatment by a healthcare professional.

